

A New Proof of the Integral Localization Formula for Equivariantly Closed Differential Forms

Matvei Libine

February 1, 2008

Abstract

In this article we give a totally new proof of the integral localization formula for equivariantly closed differential forms (Theorem 7.11 in [BGV]). We restate it here as Theorem 2.

This localization formula is very well known, but the author hopes to adapt this proof to obtain a more general result in the future.

Contents

1	Introduction	1
2	Equivariant Forms	2
3	Localization Formula	3
4	Proof of the Localization Formula	4

1 Introduction

Equivariant forms were introduced in 1950 by Henri Cartan. There are many good texts on this subject including [BGV] and [GS].

Let G be a compact Lie group acting on a compact oriented manifold M , let $\mathfrak{g}$ be the Lie algebra of G , and let $\alpha : \mathfrak{g} \rightarrow \mathcal{A}(M)$ be an equivariantly closed form on M . For $X \in \mathfrak{g}$, we denote by $M_0(X)$ the set of zeroes of the vector field on M induced by the infinitesimal action of X . We assume that $M_0(X)$ is discrete. Then Theorem 7.11 in [BGV] says that the integral

of $\alpha(X)$ can be expressed as a sum over the set of zeroes $M_0(X)$ of certain *local* quantities of M and α :

$$\int_M \alpha(X) = \sum_{p \in M_0(X)} \text{local invariant of } M \text{ and } \alpha \text{ at } p.$$

This is the essence of the integral localization formula for equivariantly closed differential forms (Theorem 2).

This localization formula for compact Lie groups G is very well known, but the problem with previously existed proofs is that they rely heavily on compactness of the acting group G and none of them generalizes to a noncompact group setting. In the future, the author intends to adapt this proof and obtain a localization formula which can be applied to some nontrivial integrals that cannot be reduced to the compact group case.

The proof which appears here is a significant modification of the localization argument which appeared in my Ph.D. thesis [L1]. This thesis contains a result on localizations of actions of noncompact groups, which is probably the only existing one on this subject. Article [L2] gives a very accessible introduction to [L1] and explains key ideas used there by way of examples.

I would like to thank Sergey Vasilyev and Michele Vergne for pointing out some very serious errors in an earlier version of this article.

2 Equivariant Forms

In this article we use the same notations as in [BGV].

Let M be a $\mathcal{C}^\infty$ -manifold of dimension n with an action of a (possibly noncompact) Lie group G , and let $\mathfrak{g}$ be the Lie algebra of G . The group G acts on $\mathcal{C}^\infty(M)$ by the formula $(g \cdot \varphi)(x) = \varphi(g^{-1}x)$. For $X \in \mathfrak{g}$, we denote by X_M the vector field on M given by (notice the minus sign)

$$(X_M \cdot \varphi)(x) = \frac{d}{d\varepsilon} \varphi(\exp(-\varepsilon X)x) \Big|_{\varepsilon=0}.$$

Let $\mathcal{A}(M)$ denote the (graded) algebra of smooth differential forms on M , and let $\mathcal{C}^\infty(\mathfrak{g}) \hat{\otimes} \mathcal{A}(M)$ denote the algebra of all smooth $\mathcal{A}(M)$ -valued functions on $\mathfrak{g}$. The group G acts on an element $\alpha \in \mathcal{C}^\infty(\mathfrak{g}) \hat{\otimes} \mathcal{A}(M)$ by the formula

$$(g \cdot \alpha)(X) = g \cdot (\alpha(g^{-1} \cdot X)) \quad \text{for all } g \in G \text{ and } X \in \mathfrak{g}.$$

Let $\mathcal{A}_G^\infty(M) = (\mathcal{C}^\infty(\mathfrak{g}) \hat{\otimes} \mathcal{A}(M))^G$ be the subalgebra of G -invariant elements. An element α of $\mathcal{A}_G^\infty(M)$ satisfies the relation $\alpha(g \cdot X) = g \cdot \alpha(X)$ and is called an *equivariant differential form*.

We define the equivariant exterior differential $d_{\mathfrak{g}}$ on $\mathcal{C}^\infty(\mathfrak{g}) \hat{\otimes} \mathcal{A}(M)$ by the formula

$$(d_{\mathfrak{g}}\alpha)(X) = d(\alpha(X)) - \iota(X_M)(\alpha(X)),$$

where $\iota(X_M)$ denotes contraction by the vector field X_M . This differential $d_{\mathfrak{g}}$ preserves $\mathcal{A}_G^\infty(M)$, and $(d_{\mathfrak{g}})^2\alpha = 0$ for all $\alpha \in \mathcal{A}_G^\infty(M)$. The elements of $\mathcal{A}_G^\infty(M)$ such that $d_{\mathfrak{g}}\alpha = 0$ are called *equivariantly closed forms*.

Example 1 Let T^*M be the cotangent bundle of M , and let σ denote the canonical symplectic form on T^*M . It is defined, for example, in [KaScha], Appendix A2. The action of the Lie group G on M naturally extends to T^*M . Then we always have a canonical equivariantly closed form on T^*M , namely, $\mu + \sigma$. Here $\mu : \mathfrak{g} \rightarrow \mathcal{C}^\infty(T^*M)$ is the moment map defined by:

$$\mu(X) : \xi \mapsto \langle \xi, X_M \rangle, \quad X \in \mathfrak{g}, \xi \in T^*M. \quad \square$$

If α is a non-homogeneous equivariant differential form, $\alpha_{[k]}$ denotes the homogeneous component of degree k . If M is a compact oriented manifold, we can integrate equivariant differential forms over M , obtaining a map

$$\int_M : \mathcal{A}_G^\infty(M) \rightarrow \mathcal{C}^\infty(\mathfrak{g})^G,$$

defined by the formula $(\int_M \alpha)(X) = \int_M \alpha(X)_{[n]}$, where $n = \dim M$.

Finally, notice that if α is an equivariantly closed form whose top homogeneous component has degree k , then $\alpha(X)_{[k]}$ is closed with respect to the ordinary exterior differential.

3 Localization Formula

From now on we assume that the Lie group G is *compact*.

We recall some more notations from [BGV]. Let $M_0(X)$ be the set of zeroes of the vector field X_M . We will state and prove the localization formula in the important special case where X_M has isolated zeroes. Here, at each point $p \in M_0(X)$, the infinitesimal action of X on $\mathcal{C}^\infty(M)$ gives rise to a linear transformation L_p on T_pM .

Because the Lie group G is compact, the transformation L_p is invertible and has only imaginary eigenvalues. Thus the dimension of M is even and

there exists an oriented basis $\{e_1, \dots, e_n\}$ of $T_p M$ such that for $1 \leq i \leq l = n/2$,

$$L_p e_{2i-1} = \lambda_{p,i} e_{2i}, \quad L_p e_{2i} = -\lambda_{p,i} e_{2i-1}.$$

We have $\det(L_p) = \lambda_{p,1}^2 \lambda_{p,2}^2 \dots \lambda_{p,l}^2$, and it is natural to take the following square root (dependent only on the orientation of the manifold):

$$\det^{1/2}(L_p) = \lambda_{p,1} \dots \lambda_{p,l}.$$

For convenience, we restate Theorem 7.11 from [BGV].

Theorem 2 *Let G be a compact Lie group with Lie algebra $\mathfrak{g}$ acting on a compact oriented manifold M , and let α be an equivariantly closed differential form on M . Let $X \in \mathfrak{g}$ be such that the vector field X_M has only isolated zeroes. Then*

$$\int_M \alpha(X) = (-2\pi)^l \sum_{p \in M_0(X)} \frac{\alpha(X)(p)}{\det^{1/2}(L_p)},$$

where $l = \dim(M)/2$, and by $\alpha(X)(p)$ we mean the value of the function $\alpha(X)_{[0]}$ at the point $p \in M$.

4 Proof of the Localization Formula

Proof of Theorem 2. Let $\pi : T^*M \rightarrow M$ denote the projection map. We regard M as a submanifold of T^*M via the zero section inclusion. Recall the canonical equivariantly closed form $\mu + \sigma$ on T^*M constructed in Example 1. We can consider the form

$$\omega(X) = e^{\mu(X) + \sigma} \wedge \pi^*(\alpha(X)), \quad X \in \mathfrak{g}. \quad (1)$$

It is an equivariantly closed form on T^*M for the reason that it is “assembled” from equivariantly closed forms. Moreover, its restriction to M is just $\alpha(X)$. We will see later that, in a way, ω is the most natural equivariant extension of α to T^*M .

Fix a G -invariant Riemannian metric on T^*M and let $\|\cdot\|$ be the norm. It induces a vector bundle diffeomorphism $TM \rightarrow T^*M$ and a norm on TM which we also denote by $\|\cdot\|$. Let $s : M \rightarrow T^*M$ denote the image of the vector field $-X_M$ under this diffeomorphism. Observe that

$$\mu(X)(s(m)) = -\|X_M\|^2 \leq 0, \quad \text{for all } m \in M,$$

and that the set

$$\{m \in M; \mu(X)(s(m)) = 0\} = \{m \in M; X_M = 0\} = M_0(X)$$

is precisely the set of zeroes of X_M .

Pick an $R > 0$ and let t increase from 0 to R . Then the family of sections $\{ts\}_{t \in [0, R]}$ provides a deformation of the initial cycle M into Rs . In other words, there exists a chain C_R in T^*M of dimension $(\dim M + 1)$ such that

$$\partial C_R = M - Rs. \quad (2)$$

Note that the support of C_R , $|C_R|$, lies inside the set

$$\{\xi \in T^*M; \mu(X)(\xi) = \langle \xi, X_M \rangle \leq 0\}. \quad (3)$$

This will ensure good behavior of the integrand when we let R tend to infinity.

Observe that if we let $R \rightarrow \infty$, then we obtain a conic Borel-Moore chain C_∞ such that

$$\partial C_\infty = M - \sum_{p \in M_0(X)} T_p^* M,$$

where each cotangent space $T_p^* M$ is provided with appropriate orientation. But we will not use this observation in the proof because one cannot interchange the order of integration and taking limit as $R \rightarrow \infty$.

Lemma 3 $(d\omega(X)_{[n]})|_{|C_R|} = 0$.

Remark 4 The form $\omega(X)_{[n]}$ itself need not be closed; this lemma only says that $\omega(X)_{[n]}$ becomes closed when restricted to $|C_R|$.

This lemma is the key moment in the proof and it is the only place where we use that the form α is equivariantly closed and that our Riemannian metric is G -invariant. Its proof is quite technical.

Proof. The form $\omega(X)_{[n]}$ can be written more explicitly as

$$\begin{aligned} \omega(X)_{[n]} &= (e^{\mu(X)+\sigma} \wedge \alpha(X))_{[n]} \\ &= e^{\mu(X)} ((e^\sigma)_{[0]} \wedge \alpha(X)_{[n]} + (e^\sigma)_{[2]} \wedge \alpha(X)_{[n-2]} + (e^\sigma)_{[4]} \wedge \alpha(X)_{[n-4]} + \dots) \\ &= e^{\mu(X)} \left(\frac{1}{0!} \sigma^0 \wedge \alpha(X)_{[n]} + \frac{1}{1!} \sigma^1 \wedge \alpha(X)_{[n-2]} + \frac{1}{2!} \sigma^2 \wedge \alpha(X)_{[n-4]} + \dots \right) \\ &= e^{\mu(X)} \sum_{k=0}^l \frac{1}{k!} \sigma^k \wedge \alpha(X)_{[n-2k]}. \end{aligned}$$

It is sufficient to prove that the form $\omega(X)_{[n]}|_{|C_R|}$ is closed away from zeroes of X_M , i.e. on the set $|C_R| \setminus M_0(X)$. So let us pick $\xi \in |C_R| \setminus M_0(X)$, say $\xi \in T_q^*M$, $q \notin M_0(X)$. Because $X_M(q) \neq 0$ we can trivialize vector field X_M near q . That is, there is a neighborhood U_q of M containing point q and positively oriented local coordinates $(x_{q,1}, \dots, x_{q,n})$ defined on U_q such that X_M is expressed by

$$X_M = \frac{\partial}{\partial x_{q,1}}. \quad (4)$$

Define a function $r : |C_R| \rightarrow \mathbb{R}$ by $r(c) = \langle c, \mu(X) \rangle$, $c \in |C_R| \subset T^*M$. Then $(x_{q,1}, \dots, x_{q,n}, r)$ defined on $\pi^{-1}(U_q) \cap |C_R|$ is a system of coordinates on $|C_R|$ containing ξ .

We expand the coordinate system $(x_{q,1}, \dots, x_{q,n})$ of M defined on U_q to the standard coordinate system $(x_{q,1}, \dots, x_{q,n}, \zeta_{q,1}, \dots, \zeta_{q,n})$ of T^*M defined on T^*U_q so that every element of T^*U_q is expressed in these coordinates as $(x_{q,1}, \dots, x_{q,n}, \zeta_{q,1}dx_{q,1} + \dots + \zeta_{q,n}dx_{q,n})$. In these coordinates,

$$\sigma = dx_{q,1} \wedge d\zeta_{q,1} + \dots + dx_{q,n} \wedge d\zeta_{q,n}.$$

Using this coordinate system $(x_{q,1}, \dots, x_{q,n}, \zeta_{q,1}, \dots, \zeta_{q,n})$ we can regard X_M as a horizontal vector field on T^*U_q . Because we chose our Riemannian metric to be G -invariant, the vector field X_M is tangent to $|C_R|$.

Using our assumption that the form α is equivariantly closed we obtain:

$$\begin{aligned} d(\omega(X)_{[n]}|_{|C_R|}) &= d\left(e^r \sum_{k=0}^l \frac{1}{k!} \sigma^k \wedge \alpha(X)_{[n-2k]}\right)\Big|_{|C_R|} \\ &= dr \wedge \left(e^r \sum_{k=0}^l \frac{1}{k!} \sigma^k \wedge \alpha(X)_{[n-2k]}\right)\Big|_{|C_R|} + e^r d\left(\sum_{k=0}^l \frac{1}{k!} \sigma^k \wedge \alpha(X)_{[n-2k]}\right)\Big|_{|C_R|} \\ &= e^r \left(\sum_{k=0}^l \frac{1}{k!} dr \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} + \sum_{k=0}^l \frac{1}{k!} \sigma^k \wedge d\alpha(X)_{[n-2k]}\right)\Big|_{|C_R|} \\ &= e^r \left(\sum_{k=0}^l \frac{1}{k!} dr \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} + \sum_{k=1}^l \frac{1}{k!} \sigma^k \wedge \iota(X_M)(\alpha(X)_{[n-2k+2]})\right)\Big|_{|C_R|} \\ &= e^r \left(\sum_{k=0}^l \frac{1}{k!} dr \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} - \sum_{k=1}^{l+1} \frac{1}{k!} \iota(X_M)(\sigma^k) \wedge \alpha(X)_{[n-2k+2]}\right)\Big|_{|C_R|} \end{aligned}$$

because $\sigma^{l+1} = 0$ and $(\sigma^k \wedge \alpha(X)_{[n-2k+2]})\Big|_{|C_R|} = 0$ for dimension reasons,

and so

$$\begin{aligned} 0 &= \iota(X_M)(\sigma^k \wedge \alpha(X)_{[n-2k+2]}) \Big|_{|C_R|} \\ &= \iota(X_M)(\sigma^k) \wedge \alpha(X)_{[n-2k+2]} \Big|_{|C_R|} + \sigma^k \wedge \iota(X_M)(\alpha(X)_{[n-2k+2]}) \Big|_{|C_R|}. \end{aligned}$$

Hence:

$$\begin{aligned} d(\omega(X)_{[n]} \Big|_{|C_R|}) &= e^r \left(\sum_{k=0}^l \frac{1}{k!} dr \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} \right. \\ &\quad \left. - \sum_{k=1}^{l+1} \frac{1}{(k-1)!} \iota(X_M)(\sigma) \wedge \sigma^{k-1} \wedge \alpha(X)_{[n-2k+2]} \right) \Big|_{|C_R|} \\ &= e^r \left(\sum_{k=0}^l \frac{1}{k!} dr \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} - \sum_{k=0}^l \frac{1}{k!} d\zeta_{q,1} \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} \right) \Big|_{|C_R|} \\ &= e^r \left(\sum_{k=0}^l \frac{1}{k!} (dr - d\zeta_{q,1}) \wedge \sigma^k \wedge \alpha(X)_{[n-2k]} \right) \Big|_{|C_R|} = 0 \end{aligned}$$

because equation (4) implies that in these coordinates, for $c \in |C_R|$,

$$\zeta_{q,1}(c) = \langle c, X_M \rangle = \langle c, \mu(X) \rangle = r(c),$$

hence $dr = d\zeta_{q,1}$. This finishes the proof of Lemma 3. $\square$

Therefore,

$$\int_M \alpha(X) = \int_M \omega(X) = \int_{Rs} \omega(X). \quad (5)$$

For each $p \in M_0(X)$, there exists a positively oriented orthonormal basis $\{e_1, \dots, e_n\}$ of $T_p M$ such that for $1 \leq i \leq l = n/2$,

$$L_p e_{2i-1} = \lambda_{p,i} e_{2i}, \quad L_p e_{2i} = -\lambda_{p,i} e_{2i-1}.$$

Using the exponential map we can find a neighborhood U_p of M containing point p and local coordinates $(x_{p,1}, \dots, x_{p,n})$ defined on U_p and centered at p such that

$$\frac{\partial}{\partial x_{p,1}}(p) = e_1, \dots, \frac{\partial}{\partial x_{p,n}}(p) = e_n$$

(in particular, the coordinate system $(x_{p,1}, \dots, x_{p,n})$ is positively oriented) and the vector field X_M is expressed by

$$X_M = \lambda_{p,1} \left(x_{p,2} \frac{\partial}{\partial x_{p,1}} - x_{p,1} \frac{\partial}{\partial x_{p,2}} \right) + \dots \\ + \lambda_{p,l} \left(x_{p,2l} \frac{\partial}{\partial x_{p,2l-1}} - x_{p,2l-1} \frac{\partial}{\partial x_{p,2l}} \right).$$

Since $M_0(X)$ is discrete, we can make these neighborhoods U_p 's small enough so that they do not overlap.

Find a sufficiently small $\varepsilon > 0$ such that, for each $p \in M_0(X)$, the cube

$$\{(t_1, \dots, t_n); |t_1| \leq \varepsilon, \dots, |t_n| \leq \varepsilon\}$$

is contained in the image of $(x_{p,1}, \dots, x_{p,n})$. For each $p \in M_0(X)$, we define a subset of M containing p

$$I_p = (x_{p,1}, \dots, x_{p,n})^{-1}(\{(t_1, \dots, t_n); |t_1| \leq \varepsilon, \dots, |t_n| \leq \varepsilon\}).$$

In formula (5) we let R tend to infinity. Recall the definition of the equivariant form ω (formula (1)). Because of the presence of the term $e^{\mu(X)+\sigma}$ there and because of the exponential decay on the set (3) we get:

$$\int_M \alpha(X) = \lim_{R \rightarrow \infty} \int_{R_s} \omega(X) = \lim_{R \rightarrow \infty} \int_{R_s \cap \bigcup_{p \in M_0(X)} \pi^{-1} I_p} \omega(X) \\ = \sum_{p \in M_0(X)} \lim_{R \rightarrow \infty} \int_{R_s \cap \pi^{-1} I_p} \omega(X).$$

That is, when we integrate over R_s , only the covectors whose basepoint lies in some I_p (i.e. "near" $M_0(X)$) count. Thus we see that the integral is localized around the set of zeroes $M_0(X)$. It remains to calculate the contribution of each point p :

$$\lim_{R \rightarrow \infty} \int_{R_s \cap \pi^{-1} I_p} \omega(X).$$

Recall that

$$\left\{ \frac{\partial}{\partial x_{p,1}} = e_1, \dots, \frac{\partial}{\partial x_{p,n}} = e_n \right\}$$

is an orthonormal basis of T_p^*M . Hence the Riemannian metric around each p can be written in terms of the frame on U_p

$$\left(\frac{\partial}{\partial x_{p,1}}, \dots, \frac{\partial}{\partial x_{p,n}} \right)$$

as $I + E_p(m)$, where I is the $n \times n$ identity matrix and $E_p(m)$ is a symmetric $n \times n$ matrix which depends on m and vanishes at the point p .

We expand each coordinate system $(x_{p,1}, \dots, x_{p,n})$ of M defined on U_p to the standard coordinate system $(x_{p,1}, \dots, x_{p,n}, \zeta_{p,1}, \dots, \zeta_{p,n})$ of T^*M defined on T^*U_p so that every element of T^*U_p is expressed in these coordinates as $(x_{p,1}, \dots, x_{p,n}, \zeta_{p,1}dx_{p,1} + \dots + \zeta_{p,n}dx_{p,n})$.

In these coordinates,

$$\begin{aligned} s = & (x_{p,1}, \dots, x_{p,n}, \lambda_{p,1}(-x_{p,2}dx_{p,1} + x_{p,1}dx_{p,2}) + \dots \\ & + \lambda_{p,l}(-x_{p,2l}dx_{p,2l-1} + x_{p,2l-1}dx_{p,2l}) + O(x^2)), \end{aligned}$$

where $O(x^2)$ denotes terms of quadratic order in $x_{p,1}, \dots, x_{p,n}$. We also have:

$$\sigma = dx_{p,1} \wedge d\zeta_{p,1} + \dots + dx_{p,n} \wedge d\zeta_{p,n},$$

and

$$\begin{aligned} \mu(X) : & (x_{p,1}, \dots, x_{p,n}, \zeta_{p,1}, \dots, \zeta_{p,n}) \\ \mapsto & \lambda_{p,1}(x_{p,2}\zeta_{p,1} - x_{p,1}\zeta_{p,2}) + \dots + \lambda_{p,l}(x_{p,2l}\zeta_{p,2l-1} - x_{p,2l-1}\zeta_{p,2l}). \end{aligned}$$

Let us choose a coordinate system for $Rs \cap \pi^{-1}I_p$, namely

$$(y_{p,1} = x_{p,1} \circ \pi, \dots, y_{p,n} = x_{p,n} \circ \pi).$$

It is positively oriented. Then in these coordinates

$$dx_{p,1} = dy_{p,1}, \dots, dx_{p,n} = dy_{p,n},$$

$$\begin{aligned} d\zeta_{p,1} = & -R\lambda_{p,1}dy_{p,2} + R \cdot O(y), & d\zeta_{p,2} = & R\lambda_{p,1}dy_{p,1} + R \cdot O(y), \dots, \\ d\zeta_{p,2l-1} = & -R\lambda_{p,l}dy_{p,2l} + R \cdot O(y), & d\zeta_{p,2l} = & R\lambda_{p,l}dy_{p,2l-1} + R \cdot O(y), \end{aligned}$$

$$\mu(X)|_{Rs} = -R\lambda_{p,1}^2(y_{p,1}^2 + y_{p,2}^2) - \dots - R\lambda_{p,l}^2(y_{p,2l-1}^2 + y_{p,2l}^2) + R \cdot O(y^3).$$

Thus $\sigma|_{Rs}$ in these coordinates becomes

$$\sigma|_{Rs} = -2R\lambda_{p,1}dy_{p,1} \wedge dy_{p,2} - \dots - 2R\lambda_{p,l}dy_{p,2l-1} \wedge dy_{p,2l} + R \cdot O(y).$$

Hence

$$(e^\sigma)_{[n]}|_{Rs} = (-2R)^l \lambda_{p,1} \dots \lambda_{p,l} dy_{p,1} \wedge \dots \wedge dy_{p,n} + R^l \cdot O(y).$$

On the other hand, the expression of $\pi^* \alpha(X)|_{Rs}$ in these coordinates does not depend on R . We have:

$$\begin{aligned} & \omega(X)_{[n]} \\ &= (e^{\mu(X)+\sigma})_{[n]} \alpha(X)_{[0]} + (e^{\mu(X)+\sigma})_{[n-2]} \alpha(X)_{[2]} + \dots + (e^{\mu(X)+\sigma})_{[0]} \alpha(X)_{[n]}. \end{aligned}$$

Hence

$$\begin{aligned} \omega(X)_{[n]}|_{Rs} &= (e^{\mu(X)+\sigma})_{[n]} \alpha(X)_{[0]}|_{Rs} \\ &+ e^{\mu(X)} \left(\text{terms depending on } R \text{ through powers of } R \text{ strictly less than } \frac{n}{2} \right) \\ &= e^{\mu(X)} (e^\sigma)_{[n]} \alpha(X)_{[0]}|_{Rs} + e^{\mu(X)} o(R^{n/2}) = \\ &e^{-R(\lambda_{p,1}^2(y_{p,1}^2+y_{p,2}^2)+\dots+\lambda_{p,l}^2(y_{p,2l-1}^2+y_{p,2l}^2)+O(y^3))} \cdot ((e^\sigma)_{[n]} \alpha(X)_{[0]}|_{Rs} + o(R^{n/2})). \end{aligned}$$

Recall that the integration takes place over I_p , hence the range of variables is $|y_{p,1}| \leq \varepsilon, \dots, |y_{p,n}| \leq \varepsilon$. We perform a change of coordinates

$$z_{p,1} = \sqrt{R}y_{p,1}, \dots, z_{p,n} = \sqrt{R}y_{p,n}, \quad |z_{p,1}| \leq \varepsilon\sqrt{R}, \dots, |z_{p,n}| \leq \varepsilon\sqrt{R}.$$

In these new coordinates,

$$\begin{aligned} \omega(X)_{[n]}|_{Rs} &= e^{-(\lambda_{p,1}^2(z_{p,1}^2+z_{p,2}^2)+\dots+\lambda_{p,l}^2(z_{p,2l-1}^2+z_{p,2l}^2)+O(z^3))/O(\sqrt{R})} \\ &\cdot ((e^\sigma)_{[n]} \alpha(X)_{[0]}|_{Rs} + 1/O(\sqrt{R})), \end{aligned}$$

and

$$(e^\sigma)_{[n]}|_{Rs} = (-2)^l \lambda_{p,1} \dots \lambda_{p,l} dz_{p,1} \wedge \dots \wedge dz_{p,n} + O(z)/O(\sqrt{R}).$$

As R tends to infinity, the integrand tends pointwise to

$$\begin{aligned} &(-2)^l \lambda_{p,1} \dots \lambda_{p,l} e^{-\lambda_{p,1}^2(z_{p,1}^2+z_{p,2}^2)-\dots-\lambda_{p,l}^2(z_{p,n-1}^2+z_{p,n}^2)} \\ &\cdot \alpha(X)_{[0]}(p) dz_{p,1} \wedge \dots \wedge dz_{p,n}. \end{aligned}$$

Hence by the Lebesgue dominated convergence theorem,

$$\begin{aligned}
\int_M \alpha(X) &= \sum_{p \in M_0(X)} \lim_{R \rightarrow \infty} \int_{Rs \cap \pi^{-1} I_p} \omega(X) \\
&= \sum_{p \in M_0(X)} \int_{\mathbb{R}^n} (-2)^l \lambda_{p,1} \dots \lambda_{p,l} \alpha(X)_{[0]}(p) \\
&\quad \cdot e^{-\lambda_{p,1}^2(z_{p,1}^2 + z_{p,2}^2) - \dots - \lambda_{p,l}^2(z_{p,n-1}^2 + z_{p,n}^2)} dz_{p,1} \wedge \dots \wedge dz_{p,n} \\
&= \sum_{p \in M_0(X)} (-2\pi)^l \frac{1}{\lambda_{p,1} \dots \lambda_{p,l}} \alpha(X)_{[0]}(p) = (-2\pi)^l \sum_{p \in M_0(X)} \frac{\alpha(X)(p)}{\det^{1/2}(L_p)}.
\end{aligned}$$

This finishes our proof of Theorem 2. $\square$

References

- [BGV] N. Berline, E. Getzler, M. Vergne, *Heat Kernels and Dirac Operators*, Springer-Verlag, 1992.
- [GS] V. Guillemin and S. Sternberg, *Supersymmetry and Equivariant de Rham Theory*, Springer-Verlag, 1999.
- [KaScha] M. Kashiwara and P. Schapira, *Sheaves on Manifolds*, Springer, 1990.
- [L1] M. Libine, *A Localization Argument for Characters of Reductive Lie Groups*, math.RT/0206019, 2002, to appear in the Jour. of Functional Analysis.
- [L2] M. Libine, *A Localization Argument for Characters of Reductive Lie Groups: An Introduction and Examples*, math.RT/0208024, 2002.

matvei@math.umass.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY
OF MASSACHUSETTS, LEDERLE GRADUATE RESEARCH TOWER,
710 NORTH PLEASANT STREET, AMHERST, MA 01003